

## Annex B

	Category 1	Category 2	Category 3
Knocked out victim's front teeth. Victim had to undergo dental treatment, and now feels reluctance to go out/nervous in the street and public places.	81.6% (169)	16.4% (34)	1.9% (4)
Spat in victim's face and beat up victim in a sustained assault resulting in a head injury (subarachnoid haemorrhage).	88.9% (184)	9.7% (20)	1.4% (3)
Bite marks to victim's arm causing pain and reddening of skin for a few days.	4.8% (10)	36.2% (75)	58.9% (122)
Injuries amounted to severe bruising and swelling and took 3 weeks to be completely healed and no longer visible.	9.2% (19)	59.9% (124)	30.9% (64)
Injuries included weakness to knee, head injury caused temporary blurred vision, and symptoms to the soft tissue of the neck caused discomfort for one week.	13.5% (28)	68.1% (141)	18.4% (38)
Deep two-inch cut to the back of victim's neck and other small cuts and scratches.	34.3% (71)	49.8% (103)	15.9% (33)
Bit victim's finger causing fracture. Significant pain and required pinning and wiring.	66.2% (137)	32.9% (68)	1.0% (2)
Pierced victim's neck with a sharp object stating it was a dirty heroin needle. Victim feared contracting HIV and/or Hepatitis B and on medication which has caused side effects making him sick. Deep distress suffered.	97.1% (201)	1.4% (3)	1.4% (3)
Injuries caused by punching and kicking, bruising from head to toe; sustained bruising to the upper left shoulder and neck, bruising to lower back, bruising to the back of right arm, bruising to the left arm, bruising to the left side of the face and neck including a black eye, bruising to legs, bruising to the side of torso.	60.9% (126)	37.2% (77)	1.9% (4)
Swollen wrist and pain to hand. Victim suffered panic attacks and needed to take 2 months off work.	42.5% (88)	54.6% (113)	2.9% (6)
Superficial cuts to neck and finger, swollen knee.	2.4% (5)	7.7% (16)	89.9% (186)
Soft tissue injuries to face; no permanent physical injury but ongoing psychological effects, waking up with nightmares and suffers anxiety.	30.9% (64)	64.3% (133)	4.8% (10)
Injuries to victim's hands causing serious discomfort but no lasting effects.	1.9% (4)	24.6% (51)	73.4% (152)
Injuries to leg; not broken but nerve damage and permanent scar, affects victim's walking and causes pain.	76.8% (159)	21.7% (45)	1.4% (3)
Multiple injuries including bruising, black eye, a bleed below the skin of the eye and a haemorrhage in inner ear.	49.3% (102)	48.8% (101)	1.9% (4)
Cuts and bruises and victim very distressed and scared to be in house alone.	22.2% (46)	66.2% (137)	11.6% (24)
Dislocated elbow, anaesthetic required to treat at hospital.	30.4% (63)	57.0% (118)	12.6% (26)
Bleeding and injury to eye, bruising and grazing to groin and leg.	9.2% (19)	61.4% (127)	29.5% (61)
Broken nose which is still deviated and several broken teeth which will require operations and medical treatment to rectify.	85.5% (177)	13.5% (28)	1.0% (2)
Pain and bruising to thighs and buttocks lasted for a week.	2.4% (5)	19.8% (41)	77.8% (161)

**Annex B**

	<b>Category 1</b>	<b>Category 2</b>	<b>Category 3</b>
Pain to head and some hair loss from hair pulling and pain to shoulder lasted a few days.	1.9% (4)	30.4% (63)	67.6% (140)
Bruising and reddening to neck from strangulation, victim feared death.	60.4% (125)	36.7% (76)	2.9% (6)
Broken nose, fully recovered after 3 weeks.	10.6% (22)	72.5% (150)	16.9% (35)
Loss of consciousness for a few minutes and temporary lump and swelling to back of head.	18.4% (38)	62.8% (130)	18.8% (39)
Victim punched 3 times in face, causing broken nose, black eyes and split lip.	37.7% (78)	53.6% (111)	8.7% (18)